

# C.U.SHAH UNIVERSITY

## Summer Examination-2018

**Subject Name : Engineering Mathematics – I**

**Subject Code : 4TE01EMT1**

**Branch: B.Tech (All)**

**Semester : 1**

**Date : 21/03/2018**

**Time : 02:30 To 05:30**

**Marks : 70**

**Instructions:**

- (1) Use of Programmable calculator & any other electronic instrument is prohibited.
- (2) Instructions written on main answer book are strictly to be obeyed.
- (3) Draw neat diagrams and figures (if necessary) at right places.
- (4) Assume suitable data if needed.

**Q-1**

**Attempt the following questions:**

**(14)**

- a) The points representing the complex number  $z$  for which  $\arg\left(\frac{z-2}{z+2}\right) = \frac{\pi}{3}$  lie on  
 (A) a circle (B) a straight line (C) an ellipse (D) a parabola
- b) If  $\left(\frac{3}{2} + \frac{i\sqrt{3}}{2}\right)^{50} = 3^{25}(x-iy)$  where  $x, y$  are real, then the ordered pair  $(x, y)$  is given by  
 (A)  $(0, 3)$  (B)  $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$  (C)  $(-3, 0)$  (D)  $(0, -3)$
- c) If  $f(x) = \frac{e^x - e^{-x}}{2}$  is continuous at  $x = 0$ , then the value of  $f(0)$  must be  
 (A) 0 (B) 1 (C) 2 (D) 3
- d)  $\lim_{x \rightarrow \infty} \frac{x^2}{e^x} = \underline{\hspace{2cm}}$   
 (A)  $-1$  (B)  $0$  (C)  $1$  (D) none of these
- e) The infinite series  $1 + r + r^2 + \dots + r^{n-1}$  is divergent if  
 (A)  $|r| < 1$  (B)  $|r| > 1$  (C)  $r \geq 1$  (D)  $r = -1$
- f) The series  $1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} + \dots \infty$  is  
 (A) convergent (B) divergent (C) absolutely convergent (D) none of these
- g) If the two tangents at the point are imaginary, the double point is called  
 (A) a node (B) a cusp (C) a conjugate point (D) none of these
- h) The curve  $x^3 + y^3 = 3axy$  represent  
 (A) Cissoid of Diocle (B) Witch of Agnesi (C) Strophoid  
 (D) Folium of Descartes



- i) The series  $x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots$  represent expansion of  
 (A)  $\sin x$  (B)  $\cos x$  (C)  $\sinh x$  (D)  $\cosh x$
- j) If  $y = \cos^{-1} x$ , then  $x$  equal to  
 (A)  $1 - \frac{y^2}{2!} + \frac{y^4}{4!} - \dots$  (B)  $y - \frac{y^3}{3!} + \frac{y^5}{5!} - \dots$  (C)  $1 + y + \frac{y^2}{2!} + \frac{y^3}{3!}$   
 (D) none of these
- k) Conditions for  $f(x, y)$  to be maximum are  
 (A)  $f_x = 0 = f_y, rt < s^2, r < 0$  (B)  $f_x = 0 = f_y, rt > s^2, r < 0$   
 (C)  $f_x = 0 = f_y, rt > s^2, r > 0$  (D)  $f_x = 0 = f_y, rt = s^2, r > 0$
- l) If  $u = y^x$ , then  $\frac{\partial u}{\partial x}$  is  
 (A)  $xy^{x-1}$  (B) 0 (C)  $y^x \log x$  (D) none of these
- m)  $\frac{\partial(u, v, w)}{\partial(x, y, z)} \cdot \frac{\partial(x, y, z)}{\partial(u, v, w)} = \text{_____}$   
 (A) 2 (B) 1 (C) 0 (D) none of these
- n) If  $f(x, y) = 0$ , then  $\frac{dy}{dx}$  is equal to  
 (A)  $\frac{\partial f / \partial x}{\partial f / \partial y}$  (B)  $\frac{\partial f / \partial y}{\partial f / \partial x}$  (C)  $-\frac{\partial f / \partial y}{\partial f / \partial x}$  (D)  $-\frac{\partial f / \partial x}{\partial f / \partial y}$

**Attempt any four questions from Q-2 to Q-8**

**Q-2 Attempt all questions (14)**

a) Prove that  $(a + ib)^{\frac{m}{n}} + (a - ib)^{\frac{m}{n}} = 2(a^2 + b^2)^{\frac{m}{2n}} \cos\left(\frac{m}{n} \tan^{-1} \frac{b}{a}\right)$ . (5)

b) Using Sandwich theorem show that (5)

(i)  $\lim_{x \rightarrow 0} x^2 \sin\left(\frac{1}{x}\right) = 0$  (ii)  $\lim_{x \rightarrow 0} x^2 \cos 20\pi x = 0$

c) Evaluate:  $\lim_{x \rightarrow 0} \frac{e^x + e^{-x} - x^2 - 2}{\sin^2 x - x^2}$  (4)

**Q-3 Attempt all questions (14)**

a) Using De Moivre's theorem prove that  $\sin 5\theta = 5\sin \theta - 20\sin^3 \theta + 16\sin^5 \theta$ . (5)

b) Prove that  $\lim_{x \rightarrow 1} \left(\frac{x}{x-1} - \frac{1}{\log x}\right) = \frac{1}{2}$ . (5)

c) Find the roots of the equation  $x^6 - i = 0$ . (4)

**Q-4 Attempt all questions (14)**

a) Expand  $f(x) = \frac{e^x}{e^x + 1}$  in powers of  $x$  up to  $x^3$  by Maclaurin's series. (5)

b) Prove that  $\tan^{-1} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$  (5)



c) Test for convergence the series  $\sum \sin\left(\frac{1}{n}\right)$ . (4)

**Q-5 Attempt all questions** (14)

a) Show that  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  is (i) convergent if  $p > 1$  and (ii) divergent if  $p \leq 1$ . (5)

b) Test for convergence the series  $\frac{1}{1 \cdot 2 \cdot 3} + \frac{x}{4 \cdot 5 \cdot 6} + \frac{x^2}{7 \cdot 8 \cdot 9} + \dots$  (5)

c) Calculate approximate value of  $\sqrt{9.12}$  by using Taylor's theorem. (4)

**Q-6 Attempt all questions** (14)

a) If  $u = \sec^{-1}\left(\frac{x^2 + y^2}{x - y}\right)$  then prove that (5)

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = -\cot u (\cot^2 u + 2).$$

b) If  $u = \log(x^3 + y^3 + z^3 - 3xyz)$  then verify  $\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z}\right)^2 u = -\frac{9}{(x + y + z)^2}$ . (5)

c) Find the asymptotes of the curve  $x^2 y^2 - x^2 y - 5xy^2 + x + y + 5 = 0$ . (4)

**Q-7 Attempt all questions** (14)

a) Trace the curve  $r^2 = a^2 \cos 2\theta$ . (5)

b) Discuss the maxima and minima of  $xy + 27\left(\frac{1}{x} + \frac{1}{y}\right)$ . (5)

c) If  $u = \tan^{-1}\left(\frac{y}{x}\right)$  then verify that  $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x}$ . (4)

**Q-8 Attempt all questions** (14)

a) Trace the curve  $y^2(2a - x) = x^3$ . (5)

b) The power consumed in an electric resistor is given by  $P = \frac{E^2}{R}$  (in watts). If  $E = 200$  volts and  $R = 8$  ohms, by how much does the power change if  $E$  is decreased by 5 volts and  $R$  is decreased by 0.20 ohms? (5)

c) If  $u = 2xy$ ,  $v = x^2 - y^2$  and  $x = r \cos \theta$ ,  $y = r \sin \theta$  then evaluate  $\frac{\partial(u, v)}{\partial(r, \theta)}$ . (4)

