Enrollment No:	Exam Seat No:
	Eaun Seut 1101

C.U.SHAH UNIVERSITY

Summer Examination-2019

Subject Name: Statistics II Subject Code: 4CO04STA2

Subject Code: 4CO04STA2

Branch: B.Com (English)

Ti 02.03 Tr. 05.20

Semester: 4 Date: 24/04/2019 Time: 02:03 To 05:30 Marks: 70

Instructions:

- (1) Use of Programmable calculator & any other electronic instrument is prohibited.
- (2) Instructions written on main answer book are strictly to be obeyed.
- (3) Draw neat diagrams and figures (if necessary) at right places.
- (4) Assume suitable data if needed.

Q-1	,	Attempt the following questions.	(14)
	a)	The normal distribution is (a) a discrete probability distribution (b) can't say	1
	b)	(c) A continuous probability distribution (d) None Standard normal variate is denoted by.	1
		(a) σ (b) μ (c) z (d) None	
	c)	The value of 'e' in normal distribution is (a) 2.7183 (b) 2.1738 (c) 3.1416 (d) 2.7138	1
	d)	What are the parameters of normal distribution?	1
		(a) x and σ (b) μ and σ (c) π and e (d) None	
	e)	In linear trend equation $y = a + bx$, a and b are known as	1
		(a) variables (b) Estimated value (c) Constants (d) Semi variables	
	f)	Trend equation $y = a + bx + cx^2$ is called	1
		(a) linear trend equation(b) Second degree parabola equation(c) Exponential trend equation(d) None of these	
	g)	Seasonal variations is a part of Variations.	1
	Ç,	(a) irregular (b) Short term (c) Cyclical (d) Long term	
	h)	Phenomena which can not be measured numerically are known as	1
	:)	(a) variable (b) Constant (c) Attributes (d) Random Variable	1
	i)	The value of Q always lies between	1
		(a) -1 and +1 (b) 0 and +1 (c) -1 and 0 (d) - ∞ and + ∞	
	j)	Co-efficient of colligation is given by	1
	k)	(a) colligation (b) Yule (c) Karl Pearson (d) Bowley All the units of the sample should be selected under conditions.	1
	K)	(a) identical (b) True (c) proper (d) None of these	1
	l)	Which is the most popular method of drawing a simple random sampling?	1
		(a) Lottery method (b) Table method	
		(c) Tossing method (d) None	
	m)	Graphics method can be used for only variable.	1
	`	(a) 3 (b) n (c) 2 (d) n-1	1
	n)	The objective function for L.P problem is	1



(a) Linear (b) Nonlinear (c) Exponential (d) None

Attempt any four questions from Q-2 to Q-8

Q-2	<i>(</i>)	Attempt all questions						(14)		
	(a)	Give the Assumption of Linear programming.							7	
	(b)	Two types of hens are kept in a poultry farm. A type of hen costs Rs. 20							7	
		each and B type of hen costs Rs. 30 each. A type of hen lays 4 eggs per week and B type of hen lays 6 eggs per week. At the most 40 hens can be								
		kept in the poultry. Not more than Rs. 1050 is to be spent on the hens. How many hens of each type should be purchased to get maximum eggs?								
Q-3		Attempt all questions (
•	(a)	Discuss the uses of Linear Programming.								
	(b)									
		are more than 64. Find mean and S.D of the distribution.								
Q-4		Attempt all questions								
	(a)) Stat the importance of normal distribution.								
	(b)									
Q-5		Attempt all							(14)	
	(a)	Give the men							7	
	(b)	Fit a $y = a + bx$ to the following time series data also compete trend								
		values & sho				_				
		X	2011	2012	201		2014	2015		
0.1		Y	15	30	48		62	85	(4.A)	
Q-6		Attempt all			.•				(14) 7	
	(a)	Explain method of studying Association. The following information is obtained from two samples drawn from two								
	(b)		-	ion is obtail	nea from	two samp	ies drawn	from two	7	
		different citie	es:			City V	(Tita V		
		Total numbe	City X City Y Total number of persons 1000 1200							
		Total number of persons 1000 1200 Literates 600 800								
		Economically backwards persons 700 700								
		Economically backwards but literates 400 400								
Q-7		Attempt all questions							(14)	
	(a)	Find the coe	-	colligation					7	
	$N = 2000$, $(A) = 260$, $(\beta) = 1720$, $(AB) = 100$									
	(b)	Stat the Adv					om sampl	ing.	7	
Q-8		Attempt all	questions						(14)	
	(a)	The families	of a localit	ty are divide	ed into tw	o strata a	ccording t	o their	7	
		economic conditions. The following information is obtained:								
		Stratum		of families		e income	Stratum			
		1		30	,	000		000		
	2 20 8,000 1,20,000									
		Find the average income of all the families of the locality. If 5								

Find the average income of all the families of the locality. If 5 families from first stratum and 3 families from second stratum are selected at random, find the variance of stratified.

(b) Form a population having observations 1,5,9,11,14 how many different sample of size three without replacement can be taken? Verify that the mean of the sample means is equal to the population mean. Also obtain the variance of the sample mean.



7