

# C.U.SHAH UNIVERSITY

## Summer Examination-2019

**Subject Name : Engineering Mathematics - 4**

**Subject Code : 4TE04EMT2**

**Branch: B.Tech (Civil/EE/Mech)**

**Semester : 4**

**Date : 15/04/2019**

**Time : 02:30 To 05:30**

**Marks : 70**

Instructions:

- (1) Use of Programmable calculator & any other electronic instrument is prohibited.
- (2) Instructions written on main answer book are strictly to be obeyed.
- (3) Draw neat diagrams and figures (if necessary) at right places.
- (4) Assume suitable data if needed.

**Q-1 Attempt the following questions:**

**(14)**

- a)  $\delta$  equal to  
 (A)  $\frac{\Delta}{E^2}$  (B)  $E^{\frac{1}{2}} + E^{-\frac{1}{2}}$  (C)  $E^{\frac{1}{2}} - E^{-\frac{1}{2}}$  (D) none of these
- b)  $E$  equal to  
 (A)  $1 + \Delta$  (B)  $\Delta \nabla$  (C)  $\nabla + \Delta$  (D)  $\nabla - \Delta$
- c) Putting  $n = 2$  in the Newton – Cote’s quadrature formula following rule is obtained  
 (A) Simpson’s  $\frac{1}{3}$  rule (B) Trapezoidal rule (C) Simpson’s  $\frac{3}{8}$  rule  
 (D) none of these
- d) While evaluating a definite integral by Trapezoidal rule, the accuracy can be increased by taking  
 (A) small number of sub – intervals (B) large number of sub – intervals  
 (C) odd number of sub – intervals (D) none of these
- e) The convergence in the Gauss – Seidel method is faster than Gauss – Jacobi method.  
 (A) True (B) False
- f) The Gauss – Jordan method in which the set of equations are transformed into diagonal matrix form.  
 (A) True (B) False
- g) Which of the following methods is the best for solving initial value problems:  
 (A) Taylor’s series method (B) Euler’s method  
 (C) Runge-Kutta method of 4<sup>th</sup> order (D) Modified Euler’s method
- h) Using modified Euler’s method, the value of  $y(0.1)$  for  $\frac{dy}{dx} = x - y$ ,  $y(0) = 1$  is  
 (A) 0.909 (B) 0.809 (C) 0.0809 (D) 0.0908
- i) The finite Fourier cosine transform of  $f(x) = 2x$ ,  $0 < x < 4$  is



- (A)  $\frac{32}{n^2 \pi^2} [(-1)^n - 1]$  (B)  $\frac{16}{n^2 \pi^2} [(-1)^n - 1]$  (C)  $\frac{32}{n^2 \pi^2} (-1)^n$  (D) none of these

j) The Fourier sine transform of  $f(x) = \begin{cases} 1, & 0 < x < a \\ 0, & x > a \end{cases}$  is

- (A)  $\sqrt{\frac{2}{\pi}} \left( \frac{1 + \cos a\lambda}{\lambda} \right)$  (B)  $\sqrt{\frac{2}{\pi}} \left( \frac{1 - \cos a\lambda}{\lambda^2} \right)$  (C)  $\sqrt{\frac{2}{\pi}} \left( \frac{1 - \cos a\lambda}{\lambda} \right)$

(D) none of these

k) If  $w = f(z) = u(x, y) + iv(x, y)$  is analytic then  $f'(z)$  equal to

- (A)  $\frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$  (B)  $\frac{\partial u}{\partial x} - i \frac{\partial v}{\partial x}$  (C)  $\frac{\partial v}{\partial y} - i \frac{\partial v}{\partial x}$  (D) none of these

l) The image of circle  $|z - 1| = 1$  in the complex plane, under the mapping  $w = \frac{1}{z}$  is

- (A)  $|w - 1| = 1$  (B)  $u^2 + v^2 = 1$  (C)  $v = \frac{1}{z}$  (D)  $u = \frac{1}{z}$

m) If  $\phi = xyz$ , the value of  $|\text{grad } \phi|$  at the point  $(1, 2, -1)$  is

- (A) 0 (B) 1 (C) 2 (D) 3

n) If  $\vec{A}(t) = 3t^2i + 4tj + 4t^3k$ ,  $\int_{t=1}^{t=2} \vec{A}(t) dt$  equal to

- (A)  $15i + 6j + 7k$  (B)  $7i + 6j + 5k$  (C)  $7i + 15j + 6k$  (D) none of these

**Attempt any four questions from Q-2 to Q-8**

**Q-2**

**Attempt all questions**

**(14)**

a) Consider following tabular values

**(5)**

x	50	100	150	200	250
y	618	724	805	906	1032

Using Newton's Backward difference interpolation formula determine  $y(300)$ .

b) Use Stirling's formula to find  $y_{28}$  given

**(5)**

that  $y_{20} = 49225$ ,  $y_{25} = 48316$ ,  $y_{30} = 47236$ ,  $y_{35} = 45926$  and  $y_{40} = 44306$ .

c) Find the finite Fourier cosine transform of  $f(x) = 2x$ ,  $0 < x < 4$ .

**(4)**

**Q-3**

**Attempt all questions**

**(14)**

a) Solve the following system of equations using Gauss-Seidel Method:

**(5)**

$30x - 2y + 3z = 75$ ,  $2x + 2y + 18z = 30$ ,  $x + 17y - 2z = 48$

b) Given that

**(5)**

x	1.00	1.05	1.10	1.15	1.20	1.25	1.30
y	1.00000	1.02470	1.04881	1.07238	1.09544	1.11803	1.14017

Find  $\frac{dy}{dx}$  at  $x = 1.05$ .

c) If  $f(z) = u + iv$  is an analytic function of  $z$  and  $u + v = e^x (\cos y + \sin y)$  then find

**(4)**

$f(z)$ .

**Q-4**

**Attempt all questions**

**(14)**

a) Apply Runge-Kutta fourth order method, to find an approximate value of  $y$  when

**(5)**



$x = 0.2$ , given that  $\frac{dy}{dx} = x + y$  and  $y = 1$  when  $x = 0$ .

b) Evaluate  $\int_0^6 \frac{1}{1+x^2} dx$  by using Simpson's 3/8<sup>th</sup> rule. (5)

c) Solve the following system of equations using Gauss-Jordan method: (4)  
 $x + 2y + z = 3$ ,  $2x + 3y + 3z = 10$ ,  $3x - y + 2z = 13$

**Q-5**

**Attempt all questions** (14)

a) Using Cauchy's integral formula, evaluate  $\oint_C \frac{z^4}{(z+1)(z-i)^2} dz$ , where C is the (5)

ellipse  $9x^2 + 4y^2 = 36$ .

b) If  $\vec{F} = (2x^2 - 4z)i - 2xyj - 8x^2k$ , then evaluate  $\iiint_V \text{div } \vec{F} dV$ , where V is (5)

bounded by the planes  $x = 0$ ,  $y = 0$ ,  $z = 0$ ,  $x + y + z = 1$ .

c) The following table gives the values of  $x$  and  $y$ : (4)

$x$	30	35	40	45	50
$y$	15.9	14.9	14.1	13.3	12.5

Use Lagrange's inverse interpolation formula to find the value of  $x$  corresponding to  $y = 13.6$ .

**Q-6**

**Attempt all questions** (14)

a) Prove that  $\vec{F} = (y^2 \cos x + z^3)i + (2y \sin x - 4)j + 3xz^2k$  is irrotational and find its (5)  
 scalar potential.

b) Under the transformation  $w = \frac{1}{z}$  (5)

(a) Find the image of  $|z - 2i| = 2$

(b) Show that the image of the hyperbola  $x^2 - y^2 = 1$  is the lemniscates

$$\rho^2 = \cos 2\theta .$$

c) Obtain Picard's second approximation solution of the initial value problem (4)

$\frac{dy}{dx} = x^2 + y^2$  for  $x = 0.4$  correct to four decimal places, given that  $y(0) = 0$ .

**Q-7**

**Attempt all questions** (14)

a) Show that the function  $f(z) = \sqrt{|xy|}$  is not analytic at the origin, although (5)  
 Cauchy-Riemann equations are satisfied.

b) Using Green's Theorem, evaluate  $\int_C (3x^2 - 8y^2)dx + (4y - 6xy)dy$  where C is the (5)

boundary of the region bounded by  $y^2 = x$  and  $y = x^2$ .

c) Evaluate  $\int_0^1 x^3 dx$  by Trapezoidal Rule using 5 subintervals. (4)

**Q-8**

**Attempt all questions** (14)

a) Solve  $\frac{dy}{dx} = x + y$  with  $y(0) = 1$  by Euler's modified method for  $x = 0.1$  correct (5)

to four decimal places by taking  $h = 0.05$ .



b) Using Fourier integral show that  $\int_0^{\infty} \frac{1 - \cos \pi \lambda}{\lambda} \sin x \lambda \, d\lambda = \begin{cases} \frac{\pi}{2} & \text{if } 0 < x < \pi \\ 0 & \text{if } x > \pi \end{cases}$  (5)

c) Prove that the angle between the surface  $x^2 + y^2 + z^2 = 9$  and  $x^2 + y^2 - z = 3$  at the point  $(2, -1, 2)$  is  $\cos^{-1}\left(\frac{8}{3\sqrt{21}}\right)$ . (4)

