

# C.U.SHAH UNIVERSITY

## Summer Examination-2019

**Subject Name : Engineering Mathematics - II**

**Subject Code : 4TE02EMT2**

**Branch: B. Tech (All)**

**Semester : 2**

**Date : 20/04/2019**

**Time : 02:30 To 05:30**

**Marks : 70**

**Instructions:**

- (1) Use of Programmable calculator & any other electronic instrument is prohibited.
- (2) Instructions written on main answer book are strictly to be obeyed.
- (3) Draw neat diagrams and figures (if necessary) at right places.
- (4) Assume suitable data if needed.

**Q-1                      Attempt the following questions:                      (14)**

a) The infinite series  $1 + r + r^2 + \dots + r^{n-1}$  is convergent if  
 (A)  $|r| < 1$  (B)  $|r| > 1$  (C)  $r = 1$  (D)  $r < -1$

b) The sum of the series  $\sum_{n=1}^{\infty} (-1)^{n-1} \frac{1}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$  is  
 (A)  $\log 2$  (B) zero (C) infinite (D) none of these

c) If  $f_n = \int_0^{\frac{\pi}{4}} \tan^n x \, dx$ , then  $(f_n + f_{n-2})$  is equal to ?  
 (A)  $\frac{1}{n}$  (B)  $\frac{1}{n-1}$  (C)  $\frac{n}{n-1}$  (D)  $\frac{n-1}{n}$

d) The value of  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^7 x \, dx$  is  
 (A)  $\frac{32\pi}{35}$  (B)  $\frac{32}{35}$  (C) zero (D)  $\frac{16}{35}$

e)  $\sqrt{\frac{1}{2}} \sqrt{\frac{3}{2}} \sqrt{\frac{5}{2}} = \text{_____}$   
 (A)  $\frac{3}{8}(\pi)^{\frac{3}{2}}$  (B)  $\frac{3}{8}(\pi)^{\frac{5}{2}}$  (C)  $\frac{3}{8}(\pi)^{\frac{1}{2}}$  (D)  $\frac{1}{8}(\pi)^{\frac{3}{2}}$

f) Duplication formula:  $\sqrt{n} \sqrt{n + \frac{1}{2}} = \text{_____}$   
 (A)  $\frac{\sqrt{\pi} \sqrt{n}}{2^{2n-1}}$  (B)  $\frac{\sqrt{\pi} \sqrt{2n}}{2^{n-1}}$  (C)  $\frac{\sqrt{\pi} \sqrt{2n}}{2^{2n-1}}$  (D)  $\frac{\sqrt{\pi} \sqrt{n}}{2^{n-1}}$

g)  $erf(x) + erf_c(x)$  is equal to  
 (A) 0 (B) 1 (C) -1 (D) 2



- h)  $\int_0^{\frac{\pi}{2}} \frac{d\theta}{\sqrt{1-2\sin^2\theta}}$  is equal to  
 (A)  $\frac{1}{\sqrt{2}}E\left(\frac{1}{\sqrt{2}}\right)$  (B)  $\frac{1}{2}K\left(\frac{1}{\sqrt{2}}\right)$  (C)  $\frac{1}{\sqrt{2}}K\left(\frac{1}{\sqrt{2}}\right)$  (D)  $\frac{1}{2}E\left(\frac{1}{\sqrt{2}}\right)$
- i) The tangents at the origin are obtained by equating to zero  
 (A) the lowest degree terms (B) the highest degree terms  
 (C) constant term (D) none of these
- j) If the powers of x are even, then the curve is symmetrical about  
 (A) X – axis (B) Y – axis (C) about both X and Y axes (D) None of these
- k)  $\int_0^{\frac{\pi}{2}} \int_0^{\infty} e^{-r^2} \cdot r \, dr \, d\theta$  is equal to  
 (A)  $\frac{\pi}{2}$  (B)  $\pi$  (C)  $\frac{\pi}{4}$  (D)  $-\frac{\pi}{4}$
- l) The transformations  $x + y = u$ ,  $x - y = v$  transform the area element  $dy \, dx$  into  $|J| \, du \, dv$ , where  $|J|$  is equal to  
 (A)  $\frac{1}{2}$  (B) 1 (C)  $u$  (D) none of these
- m) The degree and order of the differential equation of all parabolas whose axis is x-axis are  
 (A) 2, 1 (B) 1, 2 (C) 3, 2 (D) none of these
- n) Solution of differential equation  $x \, dy - y \, dx = 0$  represents  
 (A) Rectangular hyperbola (B) Circle whose centre is at origin  
 (C) Parabola whose vertex is at origin  
 (D) Straight line passing through origin

**Attempt any four questions from Q-2 to Q-8**

**Q-2 Attempt all questions (14)**

a) Using reduction formula prove that  $\int_0^a x^5 (2a^2 - x^2)^{-3} \, dx = \frac{1}{2} \left( \log 2 - \frac{1}{2} \right)$ . (5)

b) Prove that  $\int_0^{\infty} \frac{x^4}{4^x} \, dx = \frac{24}{(\log 4)^5}$ . (5)

c) Evaluate:  $\int_{-c}^c \int_{-b}^b \int_{-a}^a (x^2 + y^2 + z^2) \, dz \, dy \, dx$  (4)

**Q-3 Attempt all questions (14)**

a) Prove that  $\int_0^1 x^5 (1-x^3)^{10} \, dx = \frac{1}{3} B(2,11)$ . (5)

b) Solve:  $\frac{dy}{dx} + 2y \tan x = \sin x$  given that  $y = 0$  when  $x = \frac{\pi}{3}$  (5)



c) Test the convergence of the series  $\sum_{n=2}^{\infty} \frac{1}{n(\log n)^2}$ . (4)

**Q-4 Attempt all questions** (14)

a) By changing into polar co-ordinates, evaluate the integral (5)

$$\int_0^a \int_0^{\sqrt{a^2-y^2}} (x^2 + y^2) dx dy .$$

b) Examine the series  $1 + \frac{x}{2} + \frac{x^2}{5} + \frac{x^3}{10} + \dots + \frac{x^n}{n^2+1} + \dots$  for convergence using ratio test. (5)

c) Using reduction formula evaluate:  $\int_0^{\infty} \frac{x^4}{(1+x^2)^4} dx$  (4)

**Q-5 Attempt all questions** (14)

a) Solve:  $\frac{(x-2y) dy}{(3x+y) dx} = 3x^2 - 5xy - 2y^2$  (5)

b) Change the order of integration in the integral  $\int_0^a \int_{\frac{x^2}{a}}^{2a-x} xy dy dx$  and hence evaluate it. (5)

c) Prove that  $\int_0^{\infty} \frac{x^4(1+x^5)}{(1+x)^{15}} dx = \frac{1}{5005}$ . (4)

**Q-6 Attempt all questions** (14)

a) Examine the series  $\sum_{n=1}^{\infty} \frac{x^n}{n^p}$  for convergence using root test. (5)

b) Using reduction formula prove that  $\int_0^{\pi} x \cos^6 x dx = \frac{5\pi^2}{32}$ . (5)

c) Solve:  $(x^2 + y^2 + 1)dx - 2xy dy = 0$  (4)

**Q-7 Attempt all questions** (14)

a) Trace the curve  $y^2(2a-x) = x^3$ . (5)

b) Find the area enclosed by the cardioid  $r = a(1 - \cos \theta)$ . (5)

c) Evaluate:  $\int_0^{\frac{\pi}{2}} \frac{dx}{\sqrt{\cos x}}$  (4)

**Q-8 Attempt all questions** (14)

a) For small values of x, show that  $\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \left( x - \frac{x^3}{1!3} + \frac{x^5}{2!5} - \frac{x^7}{3!7} + \dots \right)$ . (5)

b) Trace the curve  $r = a(1 + \cos \theta)$ . (5)

c) Find the length of the arc of the curve  $y = \log \sec x$  from  $x = 0$  to  $x = \frac{\pi}{3}$  (4)

