

# C. U. SHAH UNIVERSITY

## Summer Examination-2020

**Subject Name : Engineering Mathematics – III**

**Subject Code : 4TE03EMT1**

**Branch: B. Tech (All)**

**Semester : 3**

**Date : 25/02/2020**

**Time : 02:30 To 05:30**

**Marks : 70**

**Instructions:**

- (1) Use of Programmable calculator & any other electronic instrument is prohibited.
- (2) Instructions written on main answer book are strictly to be obeyed.
- (3) Draw neat diagrams and figures (if necessary) at right places.
- (4) Assume suitable data if needed.

**Q-1**

**Attempt the following questions:**

**(14)**

- a) One of the Dirichlet's condition is function  $f(x)$  should be  
(A) single valued (B) multi valued (C) real valued (D) none of these
- b) Fourier expansion of an even function  $f(x)$  in  $(-\pi, \pi)$  has  
(A) only sine terms (B) only cosine terms  
(C) both sine and cosine terms (D) none of these
- c) In the Fourier series expansion of  $f(x) = |x|$  in  $(-\pi, \pi)$ , the value of  $b_n$  equal to  
(A) 0 (B)  $\pi$  (C)  $2\pi$  (D)  $\frac{\pi}{2}$
- d) Laplace transform of  $t^2 e^{-3t}$  is  
(A)  $\frac{\sqrt{2}}{(s+3)^2}$  (B)  $\frac{3!}{(s+3)^2}$  (C)  $\frac{2!}{(s+3)^2}$  (D)  $\frac{2!}{(s+3)^3}$
- e) Laplace transform of  $\frac{\sin t}{t}$  is  
(A)  $\cot^{-1} \frac{1}{s}$  (B)  $\tan^{-1} s$  (C)  $\tan^{-1} \frac{1}{s}$  (D)  $\sin^{-1} s$
- f)  $L^{-1} \left[ \frac{s}{(s^2 + a^2)^2} \right]$  is  
(A)  $\frac{t \cos at}{2a}$  (B)  $\frac{t^2 \sin at}{2a}$  (C)  $\frac{1}{2a^3} (\sin at - at \cos at)$  (D)  $\frac{t \sin at}{2a}$
- g)  $\frac{1}{(D-2)(D-3)(D-4)} (e^{4x} + e^{2x})$  equal to  
(A)  $4x(e^{2x} + e^{4x})$  (B)  $2(e^{2x} + e^{4x})$  (C)  $2x(e^{2x} + e^{4x})$  (D) none of these
- h) The P. I. of  $(D^2 + 1)y = \cosh 3x$  is  
(A)  $\frac{1}{10} \cosh 3x$  (B)  $\frac{1}{10} \sinh 3x$  (C)  $\frac{1}{5} \cosh 3x$  (D) none of these
- i) The C.F. of the differential equation  $(D^3 + 2D^2 + D)y = x^2$  is



- (A)  $y = c_1 + (c_2x + c_3)e^{2x}$  (B)  $y = c_1 + (c_2 + c_3x)e^{-x}$  (C)  $y = c_1 + (c_2x + c_3)e^x$   
 (D) none of these
- j) The general solution of the equation  $z = px + qy + p^2q^2$  is  
 (A)  $z = ax + by + c$  (B)  $z = ax + by + a^2 + b^2$  (C)  $z = ax + by - a^2b^2$   
 (D)  $z = ax + by + a^2b^2$
- k) The solution of the differential equation  $(1+y)p + (1+x)q = z$  is  
 (A)  $F\left(x(y-z), \frac{x+y+z}{2}\right) = 0$  (B)  $F\left(y(x-z), \frac{x+y+z}{2}\right) = 0$   
 (C)  $F\left(z(y-x), \frac{x+y+z}{2}\right) = 0$  (D) None of these
- l) The solution of  $\frac{\partial^2 z}{\partial x^2} - \frac{\partial^2 z}{\partial y^2} = 0$  is  
 (A)  $z = f_1(y+x) + f_1(y-x)$  (B)  $z = f_1(y+x) + f_2(y-x)$   
 (C)  $z = f_2(y+x) + f_2(y-x)$  (D)  $z = f(x^2 - y^2)$
- m) The order of convergence in Newton-Raphson method is  
 (A) 2 (B) 3 (C) 0 (D) None of these
- n) The order of convergence in Bisection method is  
 (A) zero (B) linear (C) quadratic (D) None of these

**Attempt any four questions from Q-2 to Q-8**

- Q-2 Attempt all questions (14)**
- a) Perform the five iteration of the Bisection method to obtain a root of the equation (5)  
 $f(x) = \cos x - xe^x$ .
- b) Find the root of the equation  $\cos x - 3x + 1 = 0$  correct to three decimal positions (5)  
 using False position method.
- c) Evaluate:  $L(t e^{2t} \cos 3t)$  (4)
- Q-3 Attempt all questions (14)**
- a) Expand  $f(x)$  in Fourier series in the interval  $(0, 2\pi)$  if (5)  
 $f(x) = \begin{cases} -\pi, & 0 < x < \pi \\ x - \pi, & \pi < x < 2\pi \end{cases}$  and show that  $\sum_{r=0}^{\infty} \frac{1}{(2r+1)^2} = \frac{\pi^2}{8}$ .
- b) Obtain Fourier series for the function  $f(x) = \begin{cases} \pi x, & 0 \leq x \leq 1 \\ \pi(2-x), & 1 \leq x \leq 2 \end{cases}$  (5)
- c) Given that one root of the equation  $x^3 - 4x + 1 = 0$  lies between 1 and 2. Find the root correct to 3 significant digits using Secant method. (4)
- Q-4 Attempt all questions (14)**
- a) Using Laplace transform method solve: (5)  
 $\frac{d^4 y}{dt^4} - k^4 y = 0, \quad y(0) = y'(0) = y''(0) = 0, y'''(0) = 1, (k \neq 0)$



b) Using convolution theorem, evaluate  $L^{-1} \left\{ \frac{s}{(s^2 + 4)^2} \right\}$ . (5)

c) Solve:  $pz - qz = z^2 + (x + y)^2$  (4)

**Q-5**

**Attempt all questions** (14)

a) Evaluate:  $L^{-1} \left[ \frac{s+2}{(s+3)(s+1)^3} \right]$  (5)

b) Solve:  $(D^2 + 5D + 4)y = x^2 + 7x + 9$  (5)

c) Solve:  $2 \frac{\partial^2 z}{\partial x^2} - 5 \frac{\partial^2 z}{\partial x \partial y} + 2 \frac{\partial^2 z}{\partial y^2} = 5 \sin(2x + y)$  (4)

**Q-6**

**Attempt all questions** (14)

a) Solve:  $(D^2 - 1)y = \cosh x \cos x$  (5)

b) Show that  $x^2 = \frac{\pi^2}{3} + 4 \sum_{n=1}^{\infty} (-1)^n \frac{\cos nx}{n^2}$  in the interval  $-\pi \leq x \leq \pi$ . Hence deduce (5)

that  $\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots = \frac{\pi^2}{12}$ .

c) Solve:  $L \left( \frac{e^{-at} - e^{-bt}}{t} \right)$  (4)

**Q-7**

**Attempt all questions** (14)

a) Solve by the method of variation of parameters:  $\frac{d^2 y}{dx^2} + a^2 y = \sec ax$  (5)

b) Solve:  $x^3 \frac{d^3 y}{dx^3} + 2x^2 \frac{d^2 y}{dx^2} + 2y = 10 \left( x + \frac{1}{x} \right)$  (5)

c) Solve:  $\frac{\partial^2 z}{\partial x^2} - 4 \frac{\partial^2 z}{\partial y^2} = \cos 2x \cos 2y$  (4)

**Q-8**

**Attempt all questions** (14)

a) Using the method of separation of variables, solve  $\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} + u$ , given (7)

$u(x, 0) = 6e^{-3x}$

b) The following table gives the variations of periodic current  $i = f(t)$  amperes over a period T sec. (7)

$t$ (sec) :	0	$\frac{T}{6}$	$\frac{T}{3}$	$\frac{T}{2}$	$\frac{2T}{3}$	$\frac{5T}{6}$	T
$i$ (A) :	1.98	1.30	1.05	1.30	-0.88	-0.5	1.98

Show, by harmonic analysis, that there is a direct current part of 0.75 amp. in the variable current and obtain the amplitude of the first harmonic.

