

C. U. SHAH UNIVERSITY

Summer Examination-2022

Subject Name: Engineering Mathematics- 4

Subject Code: 4TE04EMT2

Branch: B.Tech (Electrical)

Semester: 4

Date: 07/05/2022

Time: 11:00 To 02:00

Marks: 70

Instructions:

- (1) Use of Programmable calculator & any other electronic instrument is prohibited.
- (2) Instructions written on main answer book are strictly to be obeyed.
- (3) Draw neat diagrams and figures (if necessary) at right places.
- (4) Assume suitable data if needed.

Q-1 Attempt the following questions: (14)

- a)** $\text{curl}(\text{grad } \phi) = \underline{\hspace{2cm}}$.
- (a). 2 (b). 1 (c). 0 (d). -1
- b)** Let $f(x, y, z) = c$ represent the equation of a surface, Unit normal vector is ____
- (a). $\frac{\text{grad } f}{|\text{grad } f|}$ (b). $\text{grad } f$ (c). $\text{div}(\text{grad } f)$ (d). $\text{curl}(\text{grad } f)$
- c)** The fixed points of the mapping $w = \frac{-z+1}{z-1}$
- (a). 2, 1 (b). 1, -1 (c). 1, -2 (d). 2, -2
- d)** If $f(z) = u(x, y) + i v(x, y)$ is analytic then $f'(z) = \underline{\hspace{2cm}}$.
- (a). $u_x + i v_x$ (b). $u_x - i v_x$ (c). $u_y + i v_y$ (d). $u_x + i v_y$
- e)** The value of $\int_C \frac{dz}{z-10}$. $C: |z| = 1$
- (a). $2\pi i$ (b). $-2\pi i$ (c). $4\pi i$ (d). 0
- f)** if $\bar{A}(t) = 3t^2\hat{i} + 2t\hat{j} + 4t^3\hat{k}$, $\int_{t=1}^{t=2} \bar{A}(t)dt$ equal to
- (a). $7\hat{i} - 3\hat{j} - 5\hat{k}$ (b). $7\hat{i} + 3\hat{j} + 15\hat{k}$ (c). $-7\hat{i} - 3\hat{j} + 15\hat{k}$ (d). None of these
- g)** In Gauss- elimination method coefficient matrix reduce into
- (a). Upper triangular matrix (b). Lower triangular matrix
(c). Unit Matrix (d). Diagonal Matrix
- h)** Relation between E and Δ
- (a). $\Delta = E - 1$ (b). $\Delta = E + 1$ (c). $\Delta = 1 - E^{-1}$ (d). All of these



- i) $E^3 f(x) = \underline{\hspace{2cm}}$
 (a). $3f(x + h)$ (b). $f(x + 3h)$ (c). $f(x - 3h)$ (d). None of these
- j) Putting $n = 1$ in Newton- cote's formulae, we get _____
 (a). Trepezoidal Formula (b). Simpson's $\frac{1}{3}$ rule
 (c). Simpson's $\frac{1}{3}$ rule (d). None of these
- k) Which one of the following method is more rapid in convergence than Gauss-Jacobi method
 (a). Gauss- elimination method (b). Gauss- Jordan method
 (c). Gauss Seidel method (d). None of these
- l) If $f(x)$ is even then
 (a). $B(\lambda) = 0$ (b). $A(\lambda) = 0$ (c). Both a and b (d). None of these
- m) Write Heat Equation.
- n) Write Fourier sine integral formula.

Attempt any four questions from Q-2 to Q-8

Q-2 Attempt all questions (14)

- a) Given $\vec{u} = xyz \hat{i} + (2x^2z - y^2x) \hat{j} + xz^3 \hat{k}$ and
 (06)
 $v = xy + yz + z^2$ then find $\nabla \cdot \vec{u}$ and $\nabla \cdot v$ and $\nabla \times \vec{u}$.
- b) If $\phi(x, y, z) = x^3y + xy^3 + zxy$ then find $\nabla\phi$ and unit normal at (1,1,1). (04)
- c) Prove that $\vec{f} = \frac{x \hat{i} + y \hat{j}}{x^2 + y^2}$ is solenoidal. (04)

Q-3 Attempt all questions (14)

- a) Evaluate: $\oint_C \frac{\cos \pi z^2}{(z-1)(z-2)} dz$, where C is the circle $|z| = 3$. (05)
- b) Evaluate $\oint_C \frac{dz}{z^2 + 9}$, where C is $|z - 3i| = 4$. (05)
- c) Determine the mobius transformation that maps $z_1 = 0, z_2 = 1, z_3 = \infty$ onto $w_1 = -5, w_2 = -1, w_3 = 3$ respectively. What are the invariant points of the transformation? (04)

Q-4 Attempt all questions (14)

- a) Using Green's theorem, evaluate $\oint_C (3x^2 - 8y^2)dx + (4y - 6xy)dy$ (05)
 where C is the boundary of the region bounded by $y^2 = x$ and $y = x^2$.
- b) Verify Stoke's theorem for $\vec{F} = xy^2 \hat{i} + y \hat{j} + z^2x \hat{k}$ for the surface of a (05)
 rectangular lamina bounded by $x = 0, y = 0, x = 1, y = 2, z = 0$.
- c) Dividing the range into 10 equal parts, find the approximate value of (04)
 $\int_0^\pi \sin x dx$ by using simpson's $\frac{1}{3}$ rule.

Q-5 Attempt all questions (14)

- a) Determine the analytic function whose real part is $e^{2x}(x \cos 2y - y \sin 2y)$ (05)
- b) Solve by using Gauss-Jordan method (05)



- $2x + y + 4z = 12, \quad 8x - 3y + 2z = 20, \quad 4x + 11y - z = 33$
 c) Use Lagrange's interpolation formula to find the value of y when $x = 10$. (04)

x	5	6	9	11
y	12	13	14	16

Q-6 Attempt all questions (14)

- a) Solve the following system by using Gauss-Seidel method (05)
 $27x + 6y - z = 85, \quad 6x + 5y + 2z = 72, \quad x + y + 54z = 110$
 b) Using Taylor series method, find $y(1.1)$ correct to four decimal places, given that (05)

$$\frac{dy}{dx} = xy^{\frac{1}{3}}, y(1) = 1.$$

- c) Obtain Picard's second approximation solution of the initial value problem (04)
 $\frac{dy}{dx} = x^2 + y^2$ for $x = 0.4$ correct to four decimal places, gives that $Y(0) = 0$

Q-7 Attempt all questions (14)

- a) Given $\sin 45^\circ = 0.7071, \sin 50^\circ = 0.7660, \sin 55^\circ = 0.8192, \sin 60^\circ = 0.8660$ then find $\sin 52^\circ$ using Newton's forward Interpolation formula. (05)
 b) Find the fourier transform of $e^{-a|x|}, a > 0$ and deduce that (05)

$$\int_0^{\infty} \frac{\cos \lambda x}{\lambda^2 + a^2} d\lambda = \frac{\pi}{2a} e^{-a|x|}$$

- c) Find the fourier cosine and sine transforms of the function (04)

$$f(x) = \begin{cases} k & \text{if } 0 < x < a \\ 0 & \text{if } x > a \end{cases}$$

Q-8 Attempt all questions (14)

- a) Find the fourier transform of $e^{-(a^2x^2)}, a > 0$ and deduce that (05)

$$F\left(e^{-\frac{\lambda^2}{2}}\right) = e^{-\frac{\lambda^2}{2}}.$$

- b) Show that the function $f(z) = \sqrt{|xy|}$ is not analytic at the origin, although Cauchy-Riemann equations are satisfied. (05)
 c) If $y_0 = 3, y_1 = 12, y_2 = 81, y_3 = 2000,$ and $y_4 = 100$ then find $\Delta^4 y_0$. Also write Newton forward interpolation formula. (04)

